




Time-Limited Visual Resolution in Pilot Trainees

~~(Reprint)~~



By

19951103 032

Jeff Rabin

Aircrew Health and Performance Division

September 1995

Approved for public release; distribution unlimited.

U.S. Army Aeromedical Research Laboratory
Fort Rucker, Alabama 36362-0577

Notice

Qualified requesters

Qualified requesters may obtain copies from the Defense Technical Information Center (DTIC), Cameron Station, Alexandria, Virginia 22314. Orders will be expedited if placed through the librarian or other person designated to request documents from DTIC.

Change of address

Organizations receiving reports from the U.S. Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.

Disposition

Destroy this document when it is no longer needed. Do not return it to the originator.

Disclaimer

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citation of trade names in this report does not constitute an official Department of the Army endorsement or approval of the use of such commercial items.

Human use

Human subjects participated in these studies after giving their free and informed voluntary consent. Investigators adhered to AR 70-25 and USAMRDC Reg 70-25 on Use of Volunteers in Research.

Reviewed:




RICHARD R. LEVINE

LTC, MS


Director, Aircrew Health and
Performance Division

Released for publication:



ROGER W. WILEY, O.D., Ph.D.

Chairman, Scientific Review
Committee



DENNIS F. SHANAHAN

COL, MC, MFS
Commanding

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release, distribution unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAARL Report No.			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION U.S. Army Aeromedical Research Laboratory		6b. OFFICE SYMBOL (If applicable) MCMR-UAD	7a. NAME OF MONITORING ORGANIZATION U.S. Army Medical Research and Materiel Command		
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 620577 Fort Rucker, AL 36362-0577			7b. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, MD 21702-5012		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 0601102A	PROJECT NO. 3M161102BS15	TASK NO. PE
11. TITLE (Include Security Classification) (U) Time-Limited Visual Resolution in Pilot Trainees					
12. PERSONAL AUTHOR(S) Jeff Rabin					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day)	
15. PAGE COUNT					
16. SUPPLEMENTAL NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) visual acuity, contrast sensitivity, refractive error, visual testing		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Superior vision is needed for piloting aircraft in military and civilian environments. Although visual evaluations of potential pilots typically are conducted with no limit on viewing time, aviation and related occupations require superior vision under time-limited viewing conditions, and assessment of this capability is needed. The purpose of this study was to evaluate time-limited visual resolution in pilot trainees. A forced-choice letter recognition task was used to measure visual acuity (VA) and small letter contrast sensitivity (SLCS) in 37 trainees who had satisfied all vision requirements for pilot training. VA and SLCS were highly correlated ($r = 0.76$), indicating that the two tests measure similar aspects of visual resolution. However, although VA scores were distributed across 0.16 log units (two lines of letters on a VA chart), SLCS scores varied across 0.35 log units, which is nearly four lines on the SLCS chart. The variation in SLCS performance could be explained, in part, by subtle refractive error in pilot trainees. The results exemplify differences in performance among visually qualified trainees, and underscore the need for proper refractive correction. SLCS is a useful screening test for identifying subtle changes in vision that herald the need for optical or medical intervention.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Chief, Science Support Center			22b. TELEPHONE (Include Area Code) (334) 255-6907		22c. OFFICE SYMBOL MCMR-UAX-SI

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

Time-Limited Visual Resolution in Pilot Trainees

LTC Jeff Rabin, MS USA

Superior vision is needed for piloting aircraft in military and civilian environments. Although visual evaluations of potential pilots typically are conducted with no limit on viewing time, aviation and related occupations require superior vision under time-limited viewing conditions, and assessment of this capability is needed. The purpose of this study was to evaluate time-limited visual resolution in pilot trainees. A forced-choice letter recognition task was used to measure visual acuity (VA) and small letter contrast sensitivity (SLCS) in 37 trainees who had satisfied all vision requirements for pilot training. VA and SLCS were highly correlated ($r = 0.76$), indicating that the two tests measure similar aspects of visual resolution. However, although VA scores were distributed across 0.16 log units (two lines of letters on a VA chart), SLCS scores varied across 0.35 log units, which is nearly four lines on the SLCS chart. The variation in SLCS performance could be explained, in part, by subtle refractive error in pilot trainees. The results exemplify differences in performance among visually qualified trainees, and underscore the need for proper refractive correction. SLCS is a useful screening test for identifying subtle changes in vision that herald the need for optical or medical intervention.

Introduction

Optimal visual acuity (VA) is a requirement for piloting aircraft in the U.S. Army. Warrant officer candidates for flight school must have uncorrected VA of at least 20/20 in each eye, and no significant refractive error (-0.25 D to $+1.75$ D).¹ Implicit in these requirements is the assumption that most pilots will not require corrective lenses during flight. Spectacle wear can constrain performance when using visual aids such as helmet-mounted displays and night-vision goggles, particularly when combined with

chemical protective masks.²⁻⁴ Thus, the ideal candidate for pilot training has 20/20 vision and no obvious need for refractive correction.

But is 20/20 good enough? Most healthy young adults can see better than 20/20. In fact, 20/20 is near the lower limit of normal VA in optically corrected individuals tested under standard conditions of illumination.⁵⁻⁷ Hence, there is a range of VAs better than 20/20 that encompasses the candidates who satisfy vision requirements for pilot training. This range of vision is of interest since it may be predictive of performance differences in operational environments.

In previous studies, we showed that small letter contrast sensitivity (SLCS) is more sensitive than VA to several factors including defocus (blur),⁷ stimulus intensity,⁸ and the visual improvement achieved with two eyes compared to one.⁹ Moreover, SLCS was found to be more highly correlated than VA with small amounts of refractive error in visually normal pilot trainees.¹⁰ The value of measuring CS to assess pilot performance has been demonstrated in previous studies.^{11,12} However, SLCS is unique in its use of small letters to assess CS for recognition of fine detail. Its complements VA by providing a sensitive, adjunctive measure of visual resolution.

Although in previous studies of SLCS, and during clinical assessment of VA, observers were given ample time to recognize letter stimuli, the high speeds of aviation and other operational environments present a rapidly changing visual scene. Pilots and soldiers have limited time to process visual information in order to make critical decisions. It would be useful to assess time-limited visual resolution to determine whether it can better identify individual differences that correlate with operational performance. This paper describes time-limited VA and SLCS in pilot trainees, and how differences in SLCS can be explained, in part, by subtle refractive error, and possibly by differing capabilities that influence pilot performance.

Aircrew Health and Performance Division, U.S. Army Aeromedical Research Laboratory, Fort Rucker, AL 36362.

This manuscript was received for review in August 1994. The revised manuscript was accepted for publication in March 1995.

Methods

The stimuli for time-limited VA and SLCS were single letters generated by computer software and displayed at the center of a VGA monitor. The letters always appeared darker than the uniform, white background, which had a constant luminance of 116 cd/m². Letter contrast was controlled by software that reduced the intensity of the monitor guns relative to the white background. A cross-hair consisting of thin, black horizontal and vertical lines always appeared at the center of the screen to guide fixation to where the letter would appear periodically (but not overlapping the letter), and to help maintain the appropriate accommodative response. The entire screen subtended an angle of 3° at a viewing distance of 4.8 m.

Single letters were programmed to appear briefly (0.67 seconds) followed by an interstimulus interval of 1.66 seconds. The next letter then appeared followed by an interstimulus interval, and so on until a total of 50 letters were presented (25 VA, 25 CS), with the order of presentation randomized across VA and CS. The VA letters were of high contrast (93%) and ranged in size from 20/12.6 to 20/31.7 in 0.1 log unit steps (-0.2 to 0.2 log of the minimum angle of resolution [logMAR]).¹³ Five letters were presented at each acuity level. The SLCS letters were of constant size (20/25), but ranged in contrast from 5 to 13%, also in 0.1 log steps, with five letters per contrast step.

Testing was conducted in a darkened room with the monitor as the only source of illumination (Fig. 1). The subjects were 37 warrant officer candidates (age 21 to 29, mean = 25.6 ± 2.5 years) who previously passed all visual requirements for flight training including visual acuity, refractive error, heterophoria, and stereopsis. Subjects were tested binocularly without optical correction, since most would fly under those conditions. Each subject initially adapted to the uniform field of the monitor for about 5 minutes, during which instructions were given. Subjects were told that single letters would appear briefly at the center of the screen, and they were to verbally report each letter seen. Subjects were encouraged to give their best guess if unsure. A member of the experimental team recorded responses on a score sheet, assigning




Fig. 1. Diagram of the experimental test conditions. The observer was seated 4.8 m from the computer display on which single letters appeared briefly (0.67 seconds) centered on the screen. The letters were either high in contrast but varied, from trial to trial, in size (visual acuity), or of constant, small size (20/25) but varied in contrast (small letter contrast sensitivity).

0.02 log units per letter read correctly.^{3,5} After completing the first run (50 letters), the subject rested for 2 to 3 minutes and then completed a second run of 50 letters. All subjects gave their informed consent after protocol approval by our institutional review committee. Subjects also were assured that their responses would not be used as any form of discriminator for flight status or future training.

Results

Figure 2 shows SLCS (log contrast sensitivity) plotted against VA (logMAR) for 37 flight school candidates. Each data point is the mean of two runs corrected for the variability between runs. Correction for variability in each subject was made by subtracting one-fourth of the difference between runs from the subject's mean score. For example, a subject scoring 1.28 and 1.20 on SLCS would have a mean of 1.24, but a corrected score of $1.24 - (0.8 \div 4) = 1.22$. Yet a subject who scored 1.24 on both runs would receive the mean score of 1.24. This slight correction was used to distinguish between different levels of consistency in performance, and, as discussed subsequently, proved to be more highly correlated with subtle refractive error.

The data in Figure 2 are displayed on log scales that span equivalent ranges (0.6 log units), which facilitate direct comparison of VA and SLCS in pilot trainees. As shown in Figure 2, VA and SLCS are highly correlated ($r = 0.76$), indicating that the two tests measure similar aspects of visual resolution. However, although performance varies across 0.16 log units of VA, it is distributed across 0.36 log units of SLCS, a 60% greater range. The




Fig. 2. Small letter contrast sensitivity is plotted against visual acuity for 37 pilot trainees. Contrast sensitivity is the log of the inverse contrast threshold (logCS), and visual acuity is the log of the minimum angle of resolution (logMAR). Although logCS and logMAR are highly correlated ($r = 0.76$), scores are distributed across a more extensive range in the contrast dimension, indicating that it offers a more sensitive index of visual function. Expressed as lines of letters on a standard vision chart, scores are distributed across two lines of visual acuity, but across four lines of contrast sensitivity.

variation in VA represents only two lines of letters on a standard clinical acuity chart, whereas the variation in SLCS represents nearly four lines on the SLCS chart of similar design.^{7,10}

The greater distribution of SLCS than VA could reflect response variability unrelated to actual differences in performance. Alternatively, the larger spread of SLCS scores may indicate that SLCS is a more discriminating index of visual differences produced by small amounts of refractive error.¹⁰ To explore this issue, SLCS and VA were evaluated as a function of (cycloplegic) refractive error determined during previous flight physical examination. Since VA and SLCS were evaluated binocularly and refractive data were available for both eyes, refractive error was referenced to the most myopic meridian, which typically was the more myopic of the subject's two eyes. This was done because small amounts of myopia would have the most detrimental effect on performance,¹⁰ and preliminary analysis indicated that this approach best explained the variability in scores. Figure 3 shows mean VA (left panel) and SLCS (right panel) plotted against refractive error (lower values of logMAR indicate better VA). Although there is a tendency for VA to decrease with low myopia and higher amounts of hyperopia, this relation is not significant ($F = 3.02$, $p = 0.062$). In contrast, there is a definite peak in SLCS performance at +0.75 to +1.00 D of hyperopia, and this curvilinear relation is significant (second-order polynomial regression: $F = 8.27$, $p < 0.001$). Although refractive error explained 33% of the variability in SLCS ($r^2 = 0.33$), it accounted for only 15% of the variability in VA ($r^2 = 0.15$).

In a previous study of pilot trainees tested with unlimited viewing time, SLCS also was more highly correlated than VA with small amounts of refractive error.¹⁰ The present results extend this finding to include a larger range of refractive error and time-limited viewing conditions. Nevertheless, despite the clear correlation between SLCS and refractive error, a substantial degree of

variability in performance remains unexplained. Further research is needed to determine whether this additional variability in SLCS is predictive of differences in performance in operational environments.

Discussion

This study demonstrates that small letter contrast sensitivity provides a sensitive index of time-limited visual resolution in pilot trainees. The results are in agreement with previous studies that demonstrated the sensitivity of SLCS to defocus,⁷ luminance,⁸ binocular visual enhancement,⁹ and subtle refractive error in pilot trainees.¹⁰ Whereas in previous studies SLCS was tested with unlimited viewing time, the present results underscore the sensitivity of this approach under time-limited viewing conditions similar to what may be encountered in an operational environment.

Although VA was not strongly correlated with refractive error in the subjects tested, SLCS peaked at low hyperopia (+0.75 to +1.0 D), but declined with greater or lesser amounts of hyperopia, particularly as values approached myopia. This finding ostensibly is unclear, since subjects were uncorrected during testing, and it may be expected that best performance would occur at emmetropia (zero diopters). However, as during initial qualification for pilot status, refractive error was based on cycloplegic measurement in which accommodation (focusing ability) was neutralized pharmacologically. This eliminated any excess accommodation due to tonic factors or observer tendencies that would operate under normal viewing conditions. The consequence of neutralizing this excess accommodation is to shift refractive error toward hyperopia. Hence, a tendency to accommodate under normal (noncycloplegic) conditions in a subject with low hyperopia would optimize visual performance on SLCS, since the accommodation




Fig. 3. Mean visual acuity (logMAR) and small letter contrast sensitivity (logCS) are plotted against refractive error (lower logMAR values indicate better visual acuity). Both visual acuity and contrast sensitivity show a curvilinear relation to refractive error with peak performance between 0.75 and 1 D of hyperopia, but this relation is more prominent for contrast sensitivity.

would compensate for the refractive error. However, the emmetrope or very low myope who accommodates more than the stimulus demand would suffer a reduction in visual resolution. Although, based on cycloplegic exam, this may encourage us to select low hyperopes for flight training, a substantial degree of variability in performance remains unexplained by refractive error. Additional testing would be needed to substantiate this concept, such as measurement of the accommodative response under normal viewing conditions. The present results do indicate the importance of correcting subtle refractive error, particularly when myopic in nature. Perhaps SLCS could be implemented as a screening procedure to indicate when refractive error has changed sufficiently to warrant optical correction (Dr. Dudley Price, personal communication).

As in previous studies,⁷⁻¹⁰ the present findings underscore the greater sensitivity of SLCS than VA to factors that affect visual resolution. This is due to the precipitous, descending slope of the contrast sensitivity function (CSF). The CSF, which relates contrast sensitivity to spatial dimension, is steep near the acuity limit. Therefore, as illustrated in Figure 4, any perturbation, such as defocus, that causes a reduction in VA produces a greater decrease in SLCS. Conditions that can produce small decrements in VA, such as refractive surgery (e.g., radial keratotomy),¹⁴ edema of the cornea or retina (e.g., diabetes),¹⁵ optic neuritis (e.g., multiple sclerosis),¹⁶ and visual loss in the aging eye (e.g., cataracts),¹⁷ produce larger reductions in SLCS.¹⁸ As demonstrated in the present study, SLCS also provides a more discriminating approach for identifying candidates for occupations requiring unique visual abilities, such as those in space and aviation. Currently we are developing a hard copy, letter chart version (called




Fig. 4. The contrast sensitivity (CS) function that relates CS (inverse contrast threshold) to spatial detail (spatial frequency in cycles per degree of visual angle). The effect of defocus on the CS function is shown diagrammatically. Due to the steep descending slope of the CS function, a decrease on the x axis, which represents the loss of visual acuity (VA), is associated with a relatively larger decrease on the y axis, which represents the loss of CS.

Row		LogCS
1	U R N E D Z H F V P	0.0
2	N V Z F H E P R D U	0.1
3	D V N Z R H F U P E	0.2
4	P H V D F U E Z N R	0.3
5	R V U N D P H Z E F	0.4
6	F R E U P Z H D V R	0.5
7	E R P D N Z F U V H	0.6
USAARL Small Letter Contrast Test #1		
Row		LogCS
8	D R E Z U F V N H P	0.7
9	R P F D U N Z E V H	0.8
10	H P F E D V Z N F U	0.9
11		1.0
12		1.1
13		1.2
14		1.3
USAARL Small Letter Contrast Test #2		

Fig. 5. The newly developed Small Letter Contrast Test (SLCT), a letter chart test for general use. The SLCT has 14 lines of letters with 10 letters per line. Contrast varies by line in 0.1 log steps and credit is given for each letter read correctly (0.01 log unit per letter). Letter size is 20/25. The present study used single letters displayed on a computer monitor in time-limited fashion.

the Small Letter Contrast Test; Fig. 5) for widespread use in clinical, research, and institutional settings.¹⁸

Acknowledgments

Grateful acknowledgment is extended to SGT James Wicks and SPC Antonia Rivers for their assistance in data collection, and to Dr. Roger Wiley and LTC Richard Levine for their advice during the preparation of this manuscript.

References

- Walsh DJ, Levine RR: Compendium of US Army Visual Medical Fitness Standards. Fort Rucker, AL. US Army Aeromedical Research Laboratory, USAARL Report No. 87-11, 1987.
- Bachman WG: Contact lens wear with an aviation protective mask: a preliminary study. *Int Contact Lens Clin* 1991; 18(Jan/Feb): 21-3.
- Lattimore MJ: Contact lenses in U.S. Army attack helicopter environment: an interim report. *J Am Optom Assoc* 1992; 62: 322-5.
- Wiley RW: Military Research with Contact Lenses. Fort Rucker, AL. US Army Aeromedical Research Laboratory, USAARL Report No. 93-19, 1993.
- Sheedy JE, Bailey IL, Raasch TW: Visual acuity and chart luminance. *Am J Optom Physiol Optics* 1984; 61: 595-600.
- Pardhan S, Elliot DB: Clinical measurements of binocular summation and inhibition in patients with cataract. *Clin Vis Sci* 1991; 6: 355-9.
- Rabin J: Optical defocus: differential effects on size and contrast letter recognition thresholds. *Invest Ophthalmol Vis Sci* 1994; 35: 646-8.
- Rabin J: Two eyes are better than one: binocular enhancement in the contrast domain. *Ophthalmol Physiol Opt* 1995; 15: (in press).

9. Rabin J: Luminance effects on visual acuity and small letter contrast sensitivity. *Optom Vis Sci* 1994; 71: 685-8.
10. Rabin J: Small letter contrast sensitivity: an alternative measure of visual resolution for aviation candidates. *Aviat Space Environ Med* 1995; 66: 56-8.
11. Ginsburg AP, Evans DW, Sekuler R, et al: Contrast sensitivity predicts pilots' performance in aircraft simulators. *Am J Optom Physiol Optics* 1982; 59: 105-9.
12. Ginsburg AP, Easterly J, Evans DW: Contrast sensitivity predicts target detection field performance of pilots. In *Preprints of the Aerospace Medical Association Meeting*, pp 44-5. Alexandria, VA, Aerospace Medical Association, 1983.
13. Bailey IL, Lovie JE: New design principles for visual acuity letter charts. *Am J Optom Physiol Optics* 1976; 53: 740-5.
14. Applegate R: Acuties through annular and central pupils after radial keratotomy. *Optom Vis Sci* 1990; 68: 584-90.
15. Adams AJ: Chromatic and luminosity processing in retinal disease. *Am J Optom Physiol Optics* 1982; 5: 954-60.
16. Regan D, Silver R, Murray TJ: Visual acuity and contrast sensitivity in multiple sclerosis. *Brain* 1977; 100: 563-79.
17. Maraini G, Rosmini F, Graziosi P, et al: Influence of type and severity of pure forms of age-related cataract on visual acuity and contrast sensitivity. *Invest Ophthalmol Vis Sci* 1994; 35: 262-7.
18. Rabin J, Wicks J: Extending the range of vision assessment: the small letter contrast test. *Optom Vis Sci (Suppl)* 1994; 71: 86.